



Customer ID# 0000- 0000







Dear Customer,

Thank you for using yogi® to pursue environmental enlightenment.

Your sample tested higher than normal for mold and allergens and action is suggested, as you can see from your results here.

Live Pure's **Certified Industrial Hygienists** are available to discuss these results with you. Please schedule a **complimentary 15-minute results review**, by clicking the blue scheduler button.



- Get access to YOUR COMPLETE LAB REPORT.
- Learn more about mold exposure and health effects.
- Your results were provided by an accredited lab and reviewed by our certified industrial hygienist (CIH). To consult a CIH in your area, <u>visit the AIHA</u>
 <u>website</u> and search by zip code.

Thank you,

Live Pure Customer Service

For reminders on when to re-test your indoor air quality and drinking water and special offers as a repeat customer, join the Live Pure community today!

P.O. Box 29174 Portsmouth, NH 03802 | 603-319-8414 | <u>info@livepureinc.com</u>

© Live-Pure, Inc. 2019. Patents pending.



1501 West Knudsen Drive, Phoenix, AZ 85027 (800) 651-4802 Fax (623) 780-7695 www.emlab.com

Client: Live Pure, Inc. C/O: Customer Service

Date of Receipt: 10-30-2018 Re: 0000-0000 Date of Report: 11-01-2018

SPORE TRAP REPORT: NON-VIABLE METHODOLOGY

Location:	0 1	1:	2:						
	Outdoor Air Sample			Air Sample					
Comments (see below)	None		None						
Lab ID-Version‡:	00		00						
Analysis Date:	11/0	01/2018	11/0	01/2018					
	raw ct.	raw ct. spores/m3		spores/m3					
Ascospores	12	640	5	270					
Basidiospores	48	2,600	8	430					
Botrytis									
Chaetomium									
Cladosporium	2	110							
Curvularia									
Epicoccum			1	13					
Fusarium									
Myrothecium									
Nigrospora			1	13					
Other colorless									
Penicillium/Aspergillus types†			122	6,500					
Pithomyces				·					
Rusts									
Smuts, Periconia, Myxomycetes			7	93					
Stachybotrys									
Stemphylium									
Torula									
Ulocladium									
Zygomycetes									
Background debris (1-4+)††	1+		2+						
Hyphal fragments/m3	< 13		< 13						
Pollen/m3	< 13		< 13						
Skin cells (1-4+)	< 1+		1+						
Sample volume (liters)	75		75						
§ TOTAL SPORES/m3		3,300		7,300					

Comments:

Spore types listed without a count or data entry were not detected during the course of the analysis for the respective sample, indicating a raw

The analytical sensitivity is the spores/m³ divided by the raw count, expressed in spores/m³. The limit of detection is the analytical sensitivity (in spores/m³) multiplied by the sample volume (in liters) divided by 1000 liters.

Aerotech Laboratories, Inc

[†] The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods. Also, some species with very small spores are easily missed, and may be undercounted.

^{††}Background debris indicates the amount of non-biological particulate matter present on the trace (dust in the air) and the resulting visibility for the analyst. It is rated from 1+ (low) to 4+ (high). Counts from areas with 4+ background debris should be regarded as minimal counts and may be higher than reported. It is important to account for samples volumes when evaluating dust levels.

For more information regarding analytical sensitivity, please contact QA by calling the laboratory. \ddagger A "Version" indicated by -"x" after the Lab ID# with a value greater than 1 indicates a sample with amended data. The revision number is reflected by the value of "x".

[§] Total Spores/m3 has been rounded to two significant figures to reflect analytical precision.

1501 West Knudsen Drive, Phoenix, AZ 85027 (800) 651-4802 Fax (623) 780-7695 www.emlab.com

Client: Live Pure, Inc. C/O: Customer Service

Date of Receipt: 10-30-2018 Date of Report: 11-01-2018 Re: 0000-0000

MoldSCORETM: Spore Trap Report **Outdoor Sample:** 1 Outdoor Air Sample

Fungi Identified	Outdoor sample spores/m3						Raw	Spores/			
_	<10	0		1	K		10K	>1	00K	count	m3
Generally able to grow indoors*											
Alternaria										ND	< 13
Bipolaris/Drechslera group										ND	< 13
Chaetomium										ND	< 13
Cladosporium										2	110
Curvularia										ND	< 13
Nigrospora										ND	< 13
Penicillium/Aspergillus types†										ND	< 13
Stachybotrys										ND	< 13
Torula										ND	< 13
Seldom found growing indoors**											
Ascospores										12	640
Basidiospores										48	2,600
Rusts										ND	< 13
Smuts, Periconia, Myxomycetes										ND	< 13
Total											3,307

Location: 2 Indoor Air Sample

Fungi Identified	Indoor sample spores/m3						Raw	Spores/			
	<100)		1K			10K	>1	100I	count	m3
Generally able to grow indoors*											
Alternaria										ND	< 13
Bipolaris/Drechslera group										ND	< 13
Chaetomium										ND	< 13
Cladosporium										ND	< 13
Curvularia										ND	< 13
Epicoccum										1	13
Nigrospora										1	13
Penicillium/Aspergillus types†										122	6,500
Stachybotrys										ND	< 13
Torula										ND	< 13
Seldom found growing indoors**											
Ascospores										5	270
Basidiospores										8	430
Rusts										ND	< 13
Smuts, Periconia, Myxomycetes										7	93
Total											7,320

MoldSCORE;						
100	200	300	Score			
		ШШ	100			
			100			
			100			
			100			
			100			
			105			
			105			
			300			
			100			
			100			
			100			
			100			
			100			
			119			
Fina	l MoldSC	ORE	300			

1501 West Knudsen Drive, Phoenix, AZ 85027 (800) 651-4802 Fax (623) 780-7695 www.emlab.com

Client: Live Pure, Inc. C/O: Customer Service

C/O: Customer Service Date of Receipt: 10-30-2018 Re: 0000-0000 Date of Report: 11-01-2018

MoldSCORETM: Spore Trap Report

- * The spores in this category are generally capable of growing on wet building materials in addition to growing outdoors. Building related growth is dependent upon the fungal type, moisture level, type of material, and other factors. *Cladosporium* is one of the predominant spore types worldwide and is frequently present in high numbers. *Penicillium/Aspergillus* species colonize both outdoor and indoor wet surfaces rapidly and are very easily dispersed. Other genera are usually present in lesser numbers.
- ** These fungi are generally not found growing on wet building materials. For example, the rusts and smuts are obligate plant pathogens. However, in each group there are notable exceptions. For example, agents of wood decay are members of the basidiomycetes and high counts of a single morphological type of basidiospore on an inside sample should be considered significant.

†The spores of Aspergillus and Penicillium (and others such as Acremonium, Paecilomyces) are small and round with very few distinguishing characteristics. They cannot be differentiated by non-viable sampling methods.

‡Rated on a scale from 100 to 300. A rating less than 150 is low and indicates a low probability of spores originating inside. A rating greater than 250 is high and indicates a high probability that the spores originated from inside, presumably from indoor mold growth. A rating between 150 and 250 indicates a moderate likelihood of indoor fungal growth. MoldSCORE is NOT intended for wall cavity samples. It is intended for ambient air samples in residences. Using the analysis on other samples (like wall cavity samples) will lead to misleading results.

Aerotech Laboratories, Inc EMLab ID: 2032830, Page 2 of 2